

# Alex Psomas: Lecture 14.

Events, Conditional Probability, Independence, Bayes' Rule

1. Probability Basics Review
2. Conditional Probability
3. Independence of Events
4. Bayes' Rule

# Probability Basics Review

## Setup:

- ▶ Random Experiment.  
Flip a fair coin twice.
- ▶ Probability Space.
  - ▶ **Sample Space:** Set of outcomes,  $\Omega$ .  
 $\Omega = \{HH, HT, TH, TT\}$   
(Note: **Not**  $\Omega = \{H, T\}$  with two picks!)
  - ▶ **Probability:**  $Pr[\omega]$  for all  $\omega \in \Omega$ .  
 $Pr[HH] = \dots = Pr[TT] = 1/4$ 
    1.  $0 \leq Pr[\omega] \leq 1$ .
    2.  $\sum_{\omega \in \Omega} Pr[\omega] = 1$ .
- ▶ Event. Set of the outcomes.

# Probability is Additive

## Theorem

(a) If events  $A$  and  $B$  are disjoint, i.e.,  $A \cap B = \emptyset$ , then

$$Pr[A \cup B] = Pr[A] + Pr[B].$$

(b) If events  $A_1, \dots, A_n$  are pairwise disjoint, i.e.,  $A_k \cap A_m = \emptyset, \forall k \neq m$ , then

$$Pr[A_1 \cup \dots \cup A_n] = Pr[A_1] + \dots + Pr[A_n].$$

## Proof:

Obvious.

$$Pr[A \cup B] = \sum_{\omega \in A \cup B} Pr[\omega] = \sum_{\omega \in A} Pr[\omega] + \sum_{\omega \in B} Pr[\omega] = Pr[A] + Pr[B]$$

Can I instead say that  $|A \cup B| = |A| + |B|$ ?

**No!** We don't know if the sample space is uniform.

# Consequences of Additivity

## Theorem

(a)  $Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B];$   
(inclusion-exclusion property)

(b)  $Pr[A_1 \cup \dots \cup A_n] \leq Pr[A_1] + \dots + Pr[A_n];$   
(union bound)

(c) If  $A_1, \dots, A_N$  are a **partition** of  $\Omega$ , i.e.,  
pairwise disjoint and  $\cup_{m=1}^N A_m = \Omega$ , then

$$Pr[B] = Pr[B \cap A_1] + \dots + Pr[B \cap A_N].$$

(law of total probability)


## Proof:

(b) is obvious.

See next two slides for (a) and (c).

# Inclusion/Exclusion

$$Pr[A \cup B] = Pr[A] + Pr[B] - Pr[A \cap B]$$




$$\begin{aligned} Pr[A] &= x + y \\ Pr[B] &= y + z \\ Pr[A \cap B] &= y \\ Pr[A \cup B] &= x + y + z \end{aligned}$$

Can I instead say that  $|A \cup B| = |A| + |B| - |A \cap B|$ ?

**No!** We don't know if the sample space is uniform.

# Total probability

Assume that  $\Omega$  is the union of the disjoint sets  $A_1, \dots, A_N$ .




Then,

$$Pr[B] = Pr[A_1 \cap B] + \dots + Pr[A_N \cap B].$$

Indeed,  $B$  is the union of the disjoint sets  $A_n \cap B$  for  $n = 1, \dots, N$ .

## Roll a Red and a Blue Die.



$$|E_1 \cup E_2| = |E_1| + |E_2| - |E_1 \cap E_2|$$

$E_1$  = 'Red die shows 6';  $E_2$  = 'Blue die shows 6'

$E_1 \cup E_2$  = 'At least one die shows 6'

$$Pr[E_1] = \frac{6}{36}, Pr[E_2] = \frac{6}{36}, Pr[E_1 \cup E_2] = \frac{11}{36}.$$


## Conditional probability: example.

Two coin flips (fair coin). First flip is heads. Probability of two heads?


$\Omega = \{HH, HT, TH, TT\}$ ; Uniform probability space.

Event  $A$  = first flip is heads:  $A = \{HH, HT\}$ .

$\Omega$  : uniform



New sample space:  $A$ ; uniform still.



Event  $B$  = two heads.

The probability of two heads if the first flip is heads.

**The probability of  $B$  given  $A$  is  $1/2$ .**




## A similar example.

Two coin flips(fair coin). At least one of the flips is heads.


→ Probability of two heads?

$\Omega = \{HH, HT, TH, TT\}$ ; uniform.

Event  $A$  = at least one flip is heads.  $A = \{HH, HT, TH\}$ .



New sample space:  $A$ ; uniform still.




Event  $B$  = two heads.


The probability of two heads if at least one flip is heads.

**The probability of  $B$  given  $A$  is  $1/3$ .**

# Conditional Probability: A non-uniform example



Physical experiment



Probability model


$$\Omega = \{\text{Red, Green, Yellow, Blue}\}$$

$$Pr[\text{Red} | \text{Red or Green}] = \frac{3}{7} = \frac{Pr[\text{Red} \cap (\text{Red or Green})]}{Pr[\text{Red or Green}]}$$

## Another non-uniform example

Consider  $\Omega = \{1, 2, \dots, N\}$  with  $Pr[n] = p_n$ .

Let  $A = \{3, 4\}$ ,  $B = \{1, 2, 3\}$ .




$$Pr[A|B] = \frac{p_3}{p_1 + p_2 + p_3} = \frac{Pr[A \cap B]}{Pr[B]}.$$

## Yet another non-uniform example

Consider  $\Omega = \{1, 2, \dots, N\}$  with  $Pr[n] = p_n$ .

Let  $A = \{2, 3, 4\}$ ,  $B = \{1, 2, 3\}$ .




$$Pr[A|B] = \frac{p_2 + p_3}{p_1 + p_2 + p_3} = \frac{Pr[A \cap B]}{Pr[B]}.$$

# Conditional Probability.

**Definition:** The **conditional probability** of  $B$  given  $A$  is

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}$$



In  $A$ !


In  $B$ ?

Must be in  $A \cap B$ .

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

## More fun with conditional probability.

Toss a red and a blue die, sum is 4,  
What is probability that red is 1?




$$Pr[B|A] = \frac{|B \cap A|}{|A|} = \frac{1}{3}; \text{ versus } Pr[B] = 1/6.$$

$B$  is more likely given  $A$ .

## Yet more fun with conditional probability.

Toss a red and a blue die, sum is 7,  
what is probability that red is 1?




$$Pr[B|A] = \frac{|B \cap A|}{|A|} = \frac{1}{6}; \text{ versus } Pr[B] = \frac{1}{6}.$$

Observing  $A$  does not change your mind about the likelihood of  $B$ .

## Emptiness..

Suppose I toss 3 balls into 3 bins.

$A$  = “1st bin empty”;  $B$  = “2nd bin empty.” What is  $Pr[A|B]$ ?



$$Pr[B] = Pr[\{(a, b, c) \mid a, b, c \in \{1, 3\}\}] = Pr[\{1, 3\}^3] = \frac{8}{27}$$

$$Pr[A \cap B] = Pr[(3, 3, 3)] = \frac{1}{27}$$

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]} = \frac{(1/27)}{(8/27)} = 1/8; \text{ vs. } Pr[A] = \frac{8}{27}.$$

$A$  is less likely given  $B$ : If second bin is empty the first is more likely to have balls in it.



# Gambler's fallacy.

Flip a fair coin 51 times.

$A$  = “first 50 flips are heads”

$B$  = “the 51st is heads”

$Pr[B|A]$  ?

$A = \{HH \dots HT, HH \dots HH\}$

$B \cap A = \{HH \dots HH\}$

Uniform probability space.

$$Pr[B|A] = \frac{|B \cap A|}{|A|} = \frac{1}{2}.$$

Same as  $Pr[B]$ .

The likelihood of 51st heads does not depend on the previous flips.

# Product Rule

Recall the definition:

$$Pr[B|A] = \frac{Pr[A \cap B]}{Pr[A]}.$$

Hence,

$$Pr[A \cap B] = Pr[A]Pr[B|A].$$

Consequently,

$$\begin{aligned} Pr[A \cap B \cap C] &= Pr[(A \cap B) \cap C] \\ &= Pr[A \cap B]Pr[C|A \cap B] \\ &= Pr[A]Pr[B|A]Pr[C|A \cap B]. \end{aligned}$$

# Product Rule

## **Theorem** Product Rule

Let  $A_1, A_2, \dots, A_n$  be events. Then

$$Pr[A_1 \cap \dots \cap A_n] = Pr[A_1]Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \dots \cap A_{n-1}].$$

**Proof:** By induction.

Assume the result is true for  $n$ . (It holds for  $n = 2$ .) Then,

$$\begin{aligned} Pr[A_1 \cap \dots \cap A_n \cap A_{n+1}] \\ &= Pr[A_1 \cap \dots \cap A_n]Pr[A_{n+1}|A_1 \cap \dots \cap A_n] \\ &= Pr[A_1]Pr[A_2|A_1] \cdots Pr[A_n|A_1 \cap \dots \cap A_{n-1}]Pr[A_{n+1}|A_1 \cap \dots \cap A_n], \end{aligned}$$

so that the result holds for  $n + 1$ . □

# Correlation

An example.

Random experiment: Pick a person at random.

Event  $A$ : the person has lung cancer.

Event  $B$ : the person is a heavy smoker.

$$Pr[A|B] = 1.17 \times Pr[A].$$

Conclusion:

- ▶ Smoking increases the probability of lung cancer by 17%.
- ▶ Smoking causes lung cancer.

# Correlation

Event  $A$ : the person has lung cancer. Event  $B$ : the person is a heavy smoker.  $Pr[A|B] = 1.17 \times Pr[A]$ .

A second look.

Note that

$$\begin{aligned} Pr[A|B] = 1.17 \times Pr[A] &\Leftrightarrow \frac{Pr[A \cap B]}{Pr[B]} = 1.17 \times Pr[A] \\ &\Leftrightarrow Pr[A \cap B] = 1.17 \times Pr[A] Pr[B] \\ &\Leftrightarrow Pr[B|A] = 1.17 \times Pr[B]. \end{aligned}$$

Conclusion:

- ▶ Lung cancer increases the probability of smoking by 17%.
- ▶ Lung cancer causes smoking. Really?

## Causality vs. Correlation

Events  $A$  and  $B$  are **positively correlated** if

$$Pr[A \cap B] > Pr[A]Pr[B].$$

(E.g., smoking and lung cancer.)

$A$  and  $B$  being positively correlated does not mean that  $A$  causes  $B$  or that  $B$  causes  $A$ .

Other examples:

- ▶ Tesla owners are more likely to be rich. That does not mean that poor people should buy a Tesla to get rich.
- ▶ People who go to the opera are more likely to have a good career. That does not mean that going to the opera will improve your career.
- ▶ Rabbits eat more carrots and do not wear glasses. Are carrots good for eyesight?

# Proving Causality


Proving causality is generally difficult. One has to eliminate external causes of correlation and be able to test the cause/effect relationship (e.g., randomized clinical trials).

Some difficulties:

- ▶  $A$  and  $B$  may be positively correlated because they have a common cause. (E.g., being a rabbit.)
- ▶ If  $B$  precedes  $A$ , then  $B$  is more likely to be the cause. (E.g., smoking.) However, they could have a common cause that induces  $B$  before  $A$ . (E.g., smart, CS70, Tesla.)

# Total probability

Assume that  $\Omega$  is the union of the disjoint sets  $A_1, \dots, A_N$ .



Then,

$$Pr[B] = Pr[A_1 \cap B] + \dots + Pr[A_N \cap B].$$

Indeed,  $B$  is the union of the disjoint sets  $A_n \cap B$  for  $n = 1, \dots, N$ .


Thus,

$$Pr[B] = Pr[A_1]Pr[B|A_1] + \dots + Pr[A_N]Pr[B|A_N].$$



# Total probability

Assume that  $\Omega$  is the union of the disjoint sets  $A_1, \dots, A_N$ .



$$Pr[B] = Pr[A_1]Pr[B|A_1] + \dots + Pr[A_N]Pr[B|A_N].$$

# Independence

**Definition:** Two events  $A$  and  $B$  are **independent** if

$$Pr[A \cap B] = Pr[A]Pr[B].$$

Examples:

- ▶ When rolling two dice,  $A = \text{sum is 7}$  and  $B = \text{red die is 1}$  are independent;
- ▶ When rolling two dice,  $A = \text{sum is 3}$  and  $B = \text{red die is 1}$  are **not** independent;
- ▶ When flipping coins,  $A = \text{coin 1 yields heads}$  and  $B = \text{coin 2 yields tails}$  are independent;
- ▶ When throwing 3 balls into 3 bins,  $A = \text{bin 1 is empty}$  and  $B = \text{bin 2 is empty}$  are **not** independent;

# Independence and conditional probability

**Fact:** Two events  $A$  and  $B$  are **independent** if and only if

$$Pr[A|B] = Pr[A].$$

Indeed:  $Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$ , so that

$$Pr[A|B] = Pr[A] \Leftrightarrow \frac{Pr[A \cap B]}{Pr[B]} = Pr[A] \Leftrightarrow Pr[A \cap B] = Pr[A]Pr[B].$$

## Is you coin loaded?

Your coin is fair w.p.  $1/2$  or such that  $Pr[H] = 0.6$ , otherwise.

You flip your coin and it yields heads.

What is the probability that it is fair?

### Analysis:

$A =$  'coin is fair',  $B =$  'outcome is heads'

We want to calculate  $P[A|B]$ .

We know  $P[B|A] = 1/2$ ,  $P[B|\bar{A}] = 0.6$ ,  $Pr[A] = 1/2 = Pr[\bar{A}]$

Now,


$$\begin{aligned} Pr[B] &= Pr[A \cap B] + Pr[\bar{A} \cap B] = Pr[A]Pr[B|A] + Pr[\bar{A}]Pr[B|\bar{A}] \\ &= (1/2)(1/2) + (1/2)0.6 = 0.55. \end{aligned}$$

Thus,

$$Pr[A|B] = \frac{Pr[A]Pr[B|A]}{Pr[B]} = \frac{(1/2)(1/2)}{(1/2)(1/2) + (1/2)0.6} \approx 0.45.$$

# Is your coin loaded?

A picture:



Imagine 100 situations, among which

$m := 100(1/2)(1/2)$  are such that  $A$  and  $B$  occur and


$n := 100(1/2)(0.6)$  are such that  $\bar{A}$  and  $B$  occur.

Thus, among the  $m + n$  situations where  $B$  occurred, there are  $m$  where  $A$  occurred.

Hence,

$$Pr[A|B] = \frac{m}{m+n} = \frac{(1/2)(1/2)}{(1/2)(1/2) + (1/2)0.6}.$$

# Why do you have a fever?



Using Bayes' rule, we find


$$Pr[\text{Flu}|\text{High Fever}] = \frac{0.15 \times 0.80}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.58$$

$$Pr[\text{Ebola}|\text{High Fever}] = \frac{10^{-8} \times 1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 5 \times 10^{-8}$$

$$Pr[\text{Other}|\text{High Fever}] = \frac{0.85 \times 0.1}{0.15 \times 0.80 + 10^{-8} \times 1 + 0.85 \times 0.1} \approx 0.42$$

These are the **posterior probabilities**. One says that 'Flu' is the **Most Likely a Posteriori** (MAP) cause of the high fever.

# Bayes' Rule Operations



Bayes' Rule is the canonical example of how information changes our opinions.

# Thomas Bayes

**Thomas Bayes**




Portrait used of Bayes in a 1936 book,<sup>[1]</sup> but it is doubtful whether the portrait is actually of him.<sup>[2]</sup>

No earlier portrait or claimed portrait survives.

<b>Born</b>	c. 1701 London, England
<b>Died</b>	7 April 1761 (aged 59) <a href="#">Tunbridge Wells, Kent, England</a>
<b>Residence</b>	Tunbridge Wells, Kent, England
<b>Nationality</b>	English
<b>Known for</b>	<a href="#">Bayes' theorem</a>



# Thomas Bayes



A Bayesian picture of Thomas Bayes.

# Testing for disease.

Let's watch TV!!

Random Experiment: Pick a random male.

Outcomes: (*test, disease*)

*A* - prostate cancer.

*B* - positive PSA test.


- ▶  $Pr[A] = 0.0016$ , (.16 % of the male population is affected.)
- ▶  $Pr[B|A] = 0.80$  (80% chance of positive test with disease.)
- ▶  $Pr[B|\bar{A}] = 0.10$  (10% chance of positive test without disease.)

From [http://www.cpcn.org/01\\_psa\\_tests.htm](http://www.cpcn.org/01_psa_tests.htm) and  
<http://seer.cancer.gov/statfacts/html/prost.html> (10/12/2011.)

Positive PSA test (*B*). Do I have disease?

$$Pr[A|B]???$$

# Bayes Rule.



Using Bayes' rule, we find

$$P[A|B] = \frac{0.0016 \times 0.80}{0.0016 \times 0.80 + 0.9984 \times 0.10} = .013.$$

A 1.3% chance of prostate cancer with a positive PSA test. !!!!  
!!!!

Monty Hall.

# Summary

## Events, Conditional Probability, Independence, Bayes' Rule

### Key Ideas:

- ▶ Conditional Probability:

$$Pr[A|B] = \frac{Pr[A \cap B]}{Pr[B]}$$

- ▶ Independence:  $Pr[A \cap B] = Pr[A]Pr[B]$ .
- ▶ Bayes' Rule:

$$Pr[A_n|B] = \frac{Pr[A_n]Pr[B|A_n]}{\sum_m Pr[A_m]Pr[B|A_m]}.$$

$Pr[A_n|B]$  = posterior probability;  $Pr[A_n]$  = prior probability .

- ▶ All these are possible:

$$Pr[A|B] < Pr[A]; Pr[A|B] > Pr[A]; Pr[A|B] = Pr[A].$$